

CONTENTS

¹³ C-CP/MAS NMR studies of the cyclomalto-oligosaccharide (cyclodextrin) hydrates S.J. Heyes, N.J. Clayden and C.M. Dobson (Oxford, UK)	1
Conformational analysis of the anomeric forms of sophorose, laminarabiose, and cellobiose using MM3 M.K. Dowd (Ames, IA, USA), A.D. French (New Orleans, LA, USA) and P.J. Reilly (Ames, IA, USA)	15
The crystal and molecular structures of the pentitol penta-acetates J. Kopf, M. Morf, B. Zimmer (Hamburg, Germany) and P. Köll (Oldenburg, Germany) ..	35
Characterisation by ¹ H NMR spectroscopy of enzymically derived oligosaccharides from alkali-extractable wheat-flour arabinoxylan H. Gruppen (Wageningen, Netherlands), R.A. Hoffmann (Utrecht, Netherlands), F.J.M. Kormelink, A.G.J. Voragen (Wageningen, Netherlands), J.P. Kamerling and J.F.G. Vliegthart (Utrecht, Netherlands)	45
The application of NMR-pattern-recognition methods to the classification of peracetylated oligosaccharide residues: effects of intraclass structure D.S. Weber and W.J. Goux (Richardson, TX, USA)	65
Composition analysis of sulfoethylcelluloses by high-pH anion-exchange chromatography with pulsed amperometric detection E.A. Kragten, J.P. Kamerling, J.F.G. Vliegthart (Utrecht, Netherlands), H. Botter and J.G. Batelaan (Arnhem, Netherlands)	81
Synthesis of disialoganglioside GD1α and its positional isomer H. Prabhanjan, K. Aoyama, M. Kiso and A. Hasegawa (Gifu, Japan)	87
Syntheses of specifically deoxygenated methyl α-isomaltotriosides E. Petráková, P. Kováč and C.P.J. Glaudemans (Bethesda, MD, USA)	101
(±)-6-Acetamido-1,2-anhydro-6-deoxy- <i>myo</i> -inositol: a tight-binding inhibitor and pseudosubstrate for <i>N</i> -acetyl-β-glucosaminidases G. Legler and R. Bollhagen (Köln, FRG)	113
A spacer-modified disaccharide as a photoaffinity reagent for the acceptor-binding area of bovine (1 → 4)-β-D-galactosyltransferase: comparison of its acceptor properties with those of other 2-acetamido-2-deoxy-β-D-glucopyranosides S.-C. Áts, J. Lehmann and S. Petry (Freiburg, FRG)	125
Spacer-modified trisaccharide glycosides that mimic the biantennary Asn-linked oligosaccharide acceptor of (1 → 4)-β-D-galactosyltransferase and can be used as competitive inhibitors and for irreversible deactivation S.-C. Áts, J. Lehmann and S. Petry (Freiburg, FRG)	141

Structural characterization of a novel diglycosyl diacylglyceride glycolipid from <i>Rhizobium trifolii</i> ANU843 G.G. Orgambide, R.I. Hollingsworth and F.B. Dazzo (East Lansing, MI, USA)	151
The amylose–iodine complex K.A. Murdoch (Glasgow, UK)	161
Artificial antigens. Synthetic carbohydrate haptens immobilized on crystalline bacterial surface layer glycoproteins P. Messner, M.A. Mazid, F.M. Unger (Edmonton, AB, Canada) and U.B. Sleytr (Wien, Austria)	175
Structure of the polysaccharide chains of <i>Pseudomonas pseudomallei</i> lipopolysaccharides Y.A. Knirel, N.A. Paramonov, A.S. Shashkov, N.K. Kochetkov (Moscow, Russia), R.G. Yarullin, S.M. Farber and V.I. Efremenko (Volgograd, Russia)	185
Structural elucidation of the capsular polysaccharide of <i>E. coli</i> serotype K47 A.H. De Bruin, H. Parolis and L.A.S. Parolis (Grahamstown, South Africa)	195
Structural variability in the glucuronoxylomannan of <i>Cryptococcus neoformans</i> serotype A isolates determined by ^{13}C NMR spectroscopy S.H. Turner, R. Cherniak, E. Reiss (Atlanta, GA, USA) and K.J. Kwon-Chung (Bethesda, MD, USA)	205
<i>Notes</i>	
Estimation of dialdehyde groups in 2,3-dialdehyde bead-cellulose K. Pommerening, H. Rein (Berlin, Germany), D. Bertram and R. Müller (Leipzig, Germany)	219
Struktur von 4-(Adenin-9-yl)-1,6-anhydro-4-desoxy- β -D-glucopyranose Dihydrat P. Luger und E. Rabius (Berlin, Deutschland)	225
Negative-ion fast-atom-bombardment mass spectrometry of native gangliosides using a high-polar matrix system R. Isobe, R. Higuchi and T. Komori (Fukuoka, Japan)	231
Assignment of ^{13}C NMR signals for reduced nigerooligosaccharides prepared by partial acid hydrolysis of (1 \rightarrow 3)- α -D-glucan A. Shimamura, Y. Uezono, H. Tsumori and H. Mukasa (Saitama, Japan)	237
2,3,4,6-Tetra-O-benzyl- β -D-galactopyranosyl phenyl sulfoxide as a glycosyl donor. Synthesis of some oligosaccharides containing an α -D-galactopyranosyl group A.K. Sarkar and K.L. Matta (Buffalo, NY, USA)	245
A novel route to 2-deoxy-2-iodo-D-mannopyranose derivatives D. Miljković, E. Djurendić, N. Vukojević, K. Gaši and J. Csanádi (Novi Sad, Yugoslavia)	251
A derivative of chitosan and 2,4-pentanedione with strong chelating properties M. Gómez-Guillén, A. Gómez-Sánchez and M.-E. Martín-Zamora (Seville, Spain)	255
An improved synthesis of D-amicetose S. Lajšić, D. Miljković and G. Četković (Novi Sad, Yugoslavia)	261

A linear (1 → 5)-linked α -L-arabinofuranan from the seeds of guapuruvu (<i>Schizolobium parahy- bum</i>) S.F. Zawadzki-Baggio, M.-R. Sierakowski, J.B.C. Corrêa and F. Reicher (Curitiba, Brazil)	265
The major capsular polysaccharide of <i>Cryptococcus neoformans</i> serotype B A.K. Bhattacharjee (Washington, DC, USA), K.J. Kwon-Chung and C.P.J. Glaudemans (Bethesda, MD, USA)	271
Properties of two β -glucosidases purified from the termite <i>Macrotermes muelleri</i> and from its symbiotic fungus <i>Termitomyces</i> sp. C. Rouland, M. Matoub, P. Mora (Créteil, France) and F. Petek (Villejuif, France)	273
<i>Preliminary communications</i>	
A new and extremely fast synthesis of 2-deoxy-2,2-difluoro-D-arabino-hexose (2-deoxy-2,2-di- fluoro-D-glucose) A. El-Laghdach, M.I. Matheu, S. Castellón (Tarragona, Spain), C. Bliard, A. Olesker and G. Lukacs (Gif-sur-Yvette, France)	C1
Expedient syntheses of inososes from carbohydrates: conformational and stereoelectronic as- pects of the Ferrier reaction A.S. Machado (Gerais, Brazil), D. Dubreuil, J. Cleophax, S.D. Gero (Gif-sur-Yvette, France) and N.F. Thomas (Bath, UK)	C5
<i>Author index</i>	C9
<i>Subject index</i>	C11
<i>Contents</i>	C15

